



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\begin{aligned} \lim_{x \rightarrow 0} \left( \frac{\tan x}{x} \right)^{1/x^n} &= \lim_{x \rightarrow 0} \left\{ \left[ 1 + \frac{x^2}{3} (1 + \epsilon) \right]^{x^2 (1 + \epsilon)^{-1}} \right\}^{\frac{1 + \epsilon}{3 x^{n-2}}} = e^{\lim_{x \rightarrow 0} \frac{1 + \epsilon}{3 x^{n-2}}} \\ &= \left( \epsilon \frac{1}{3 x^{n-2}} \right)_{x=0}. \end{aligned}$$

This has the advantage of being a direct, closed expression, which shows at once the particular results,  $1, e^{\frac{1}{3}}, \infty$ , when  $n$  equals  $1, 2, 2+r$ , where  $r$  is any positive integer, and also the general law for any value of  $n$ , of whatever character it may be.

### MECHANICS.

73. Proposed by J. K. ELLWOOD, A. M., Principal of Colfax School, Pittsburg, Pa.

A sixteen foot plank weighs thirty-two pounds and is supported by two props, four feet and two feet from the ends. What weight is supported by each prop?

Solution by C. HORNUNG, A. M., Professor of Mathematics, Heidelberg University, Tiffin, Ohio; P. S. BERG, B. Sc., Principal of Schools, Larimore, N. D.; J. SCHEFFER, A. M., Hagerstown, Md.; CHARLES C. CROSS, Libertytown, Md.; ELMER SCHUYLER, High Bridge, N. J.; A. H. BELL, Hillsboro, Ill.; CHARLES E. MEYERS, Canton, O., and M. A. GRUBER, A. M., War Department, Washington, D. C.

Let  $x$  = the number of pounds supported by the prop four feet from the end. Then taking moments around the other point of support, we have:

$$(6+4)x = 6 \times 32, \text{ or } x = 19.2 \text{ pounds.}$$

The other prop supports  $32 - 19.2 = 12.8$  pounds.

Also solved by G. B. M. ZERR.

74. Proposed by B. F. FINKEL, A. M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

In the experiment of swinging in a vertical circle a glass containing water, and suspended by means of a string, if the string be two feet long, what must be the velocity at the lowest point if the experiment is to succeed? [From *Ziwet's Theoretical Mechanics*, Part III., p. 96.]

Solution by the PROPOSER.

Let  $M$  = the mass of the glass and water,  $f$  = the centrifugal force,  $r$  = 2 feet = radius. Then  $f = Mv^2/r$ .

At the highest point, in order that the experiment may be successful  $f = Mg$ .  
 $\therefore Mv^2/r = Mg$ , whence  $v = \sqrt{rg}$ .

The velocity,  $v_1$ , due to gravity in passing from the highest to the lowest point is  $v_1 = \sqrt{(2gs)}$ , where  $s = 2r$ .

Hence, the velocity,  $v_2$ , at the lowest point is,

$v_2 = \sqrt{(v^2 + v_1^2)} = \sqrt{(rg + 2gs)} = \sqrt{(321.6)} = 17.94$  feet per second,  
 $g$  being equal to 32.16.

Also solved with slightly different results by G. B. M. ZERR, and CHARLES E. MEYERS.